Applicant: Hsuan-Yin Lan-Handst et al

Serial No.: 10/025,947

Filed: December 26, 2001


Page : 2

C₁₋₄ alkoxy, hydroxyl, halo, amino, nitro, cyano, C₃₋₅ cycloalkyl, 3-5 membered heterocycloalkyl, monocyclic aryl, 5-6 membered heteroaryl, C₁₋₄ alkylcarbonyloxy, C₁₋₄ alkyloxycarbonyl, C₁₋₄ alkylcarbonyl, or formyl; and further being optionally interrupted by -O-, -N(R^c)-, -N(R^c)-C(O)-O-, -O-C(O)-N(R^c)-, -N(R^c)-C(O)-N(R^d)-, or -O-C(O)-O-; each of R^c and R^d, independently, being hydrogen, alkyl, alkenyl, alkynyl, alkoxy, hydroxylalkyl, hydroxyl, or haloalkyl; provided that when L contains two or more double bonds, the double bonds are not adjacent to each other; that when L contains three double bonds, said hydrocarbon chain is substituted with C₁₋₄ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, C₁₋₄ alkoxy, hydroxyl, halo, amino, nitro, cyano, C₃₋₅ cycloalkyl, 3-5 membered heterocycloalkyl, monocyclic aryl, 5-6 membered heteroaryl, C₁₋₄ alkylcarbonyloxy, C₁₋₄ alkyloxycarbonyl, C₁₋₄ alkylcarbonyl, or formyl; and further provided that when L contains 7 carbon atoms or fewer in the hydrocarbon chain and A is C₁₋₄ alkyl phenyl or unsubstituted phenyl, Y¹ is not a bond or CH₂;

's Docket No.: 12938-003002

or a salt thereof. A

A22. (Amended) A compound of formula (I):



wherein

A is a cyclic moiety selected from the group consisting of aryl and heteroaryl; the cyclic moiety being optionally substituted with alkyl, alkenyl, alkynyl, alkoxy, hydroxylalkyl, or amino;

each of X¹ and X², independently, is O or S;

each of Y^1 and Y^2 , independently, is $-CH_2$ -, -O-, -S-, $-N(R^a)$ -, $-N(R^a)$ -C(O)-O-, -O-C(O)- $N(R^a)$ -, $-N(R^a)$ -C(O)- $N(R^b)$ -, -O-C(O)-O-, or a bond; each of R^a and R^b , independently, being hydrogen, alkyl, hydroxyla kyl, or haloalkyl;

L is a straight C_{3-12} hydrocarbon chain optionally containing at least one double bond, at least one triple bond, or at least one double bond and one triple bond; said hydrocarbon chain

B